

77)

**Structure of (—)-Neoverrucosan-5 $\beta$ -ol, a Diterpenoid from *Mylia verrucosa*  
(Liverwort) containing a Novel Carbon Skeleton: X-Ray Crystal  
and Molecular Structure of the Benzoate**

By AKIHIKO MATSUO,\* HIROSHI NOZAKI, and MITSURU NAKAYAMA

*(Department of Chemistry, Faculty of Science, Hiroshima University, Naka-ku, Hiroshima 730, Japan)*

and DAISUKE TAKAOKA

*(Department of Chemistry, Faculty of General Education, Ehime University, Matsuyama 790, Japan)*

and SHŪICHI HAYASHI

*(Department of Chemistry, Faculty of Science, Okayama University of Science, Okayama 700, Japan)*

**Reprinted from the Journal of The Chemical Society  
Chemical Communications 1980**

## Structure of (—)-Neoverrucosan-5 $\beta$ -ol, a Diterpenoid from *Mylia verrucosa* (Liverwort) containing a Novel Carbon Skeleton: X-Ray Crystal and Molecular Structure of the Benzoate

By AKIHIKO MATSUO,\* HIROSHI NOZAKI, and MITSURU NAKAYAMA

(Department of Chemistry, Faculty of Science, Hiroshima University, Naka-ku, Hiroshima 730, Japan)

and DAISUKE TAKAOKA


(Department of Chemistry, Faculty of General Education, Ehime University, Matsuyama 790, Japan)

and SHŪICHI HAYASHI

(Department of Chemistry, Faculty of Science, Okayama University of Science, Okayama 700, Japan)


**Summary** The structure and absolute configuration of a novel diterpene alcohol, (—)-neoverrucosan-5 $\beta$ -ol, isolated from the liverwort *Mylia verrucosa* has been shown to be (2) on the basis of chemical and spectral evidence and X-ray analysis of the benzoate (6).

PREVIOUSLY, we have isolated (—)-2 $\beta$ -hydroxyverrucosane (1) and five related diterpenoids with a verrucosane skeleton from the liverwort *Mylia verrucosa* Lindb., and their structures, including absolute configuration, have been determined by extensive chemical reactions and spectral data.<sup>1-3</sup> The novel verrucosane skeleton has been recently confirmed by X-ray analysis of the mono-*p*-bromobenzoate of the major diol.<sup>4</sup> In continuing research on chemical constituents of the liverwort, we have isolated a further diterpene alcohol, (—)-neoverrucosan-5 $\beta$ -ol (2), containing a novel carbon skeleton† from the ethanolic extract by means of column chromatography and preparative t.l.c. We describe here not only the structure and absolute configuration of the novel alcohol (2), but also its acid-catalysed reaction which gives (+)-homoverrucosan-5 $\beta$ -ol (7).



The i.r. and <sup>1</sup>H n.m.r. spectra‡ of the compound (2), C<sub>20</sub>H<sub>34</sub>O, m.p. 174–175 °C, [ $\alpha$ ]<sub>D</sub> –10°, revealed that it was a saturated tetracyclic diterpenoid containing a cyclopropane ring [ $\nu$  3060 cm<sup>-1</sup>;  $\delta$  0.2–0.7 (2H, m)], an isopropyl group [ $\nu$  1380 and 1373 cm<sup>-1</sup>;  $\delta$  0.81 and 0.88 (each 3H, d,  $J$  = 5.5 Hz)], 3 tertiary methyls [ $\delta$  0.73, 0.82, and 1.17 (each 3H, s)], and a secondary hydroxy group [ $\nu$  3270 cm<sup>-1</sup>;  $\delta$  4.01 (1H, dd,  $J$  = 7.0 and 4.5 Hz)], the presence of which was confirmed by formation of the acetate (3), C<sub>22</sub>H<sub>36</sub>O<sub>2</sub>, m.p. 127.5–129 °C, [ $\alpha$ ]<sub>D</sub> –6°. Jones reagent oxidized the alcohol (2) to the ketone (4), C<sub>20</sub>H<sub>32</sub>O, m.p. 67.5–68 °C, [ $\alpha$ ]<sub>D</sub> +140°. It had a carbonyl group (6-membered ketone)

conjugated with the cyclopropane ring [ $\nu$  1670 cm<sup>-1</sup>] and a methylene adjacent to a quaternary carbon atom [ $\nu$  1435 cm<sup>-1</sup>;  $\delta$  1.87 and 2.06 (each 1H, d,  $J$  15.0 Hz)]. The ketone (4) was then reduced with NaBH<sub>4</sub> to afford the original alcohol (2) and its epimer (5), C<sub>20</sub>H<sub>34</sub>O, m.p. 121–122 °C, [ $\alpha$ ]<sub>D</sub> –52°, in a ratio of 2:5. This chemical evidence and the similarity of the spectra with those of verrucosanol (1) and its derivatives suggested that the overall structure might be represented by (2).



Reagents: i, Ac<sub>2</sub>O in pyridine; ii, CrO<sub>3</sub> in H<sub>2</sub>SO<sub>4</sub>-acetone; iii, NaBH<sub>4</sub> in EtOH; iv, PhCOCl in pyridine; v, 0.5 N H<sub>2</sub>SO<sub>4</sub> in acetone (1:5).

† We propose the name neoverrucosane for the new carbon skeleton from its connection with the verrucosane skeleton, and suggest the numbering shown in (2).

‡ I.r. and <sup>1</sup>H n.m.r. spectra were, respectively, determined for KBr pellets and CDCl<sub>3</sub> solution, and optical rotations for CHCl<sub>3</sub> solution. All new compounds (2)–(6) gave spectral data in good agreement with the assigned structures.

In order to confirm the structure and stereochemistry, the alcohol (2) was transformed into the benzoate (6),  $C_{27}H_{36}O_2$ , m.p. 148.5–150 °C,  $[\alpha]_D +13^\circ$ , colourless orthorhombic crystals, and an X-ray analysis of the benzoate was carried out.




FIGURE. A computer-generated perspective drawing of the benzoate (6).

Crystal data:  $a = 6.278(2)$ ,  $b = 9.971(6)$ ,  $c = 37.451(26)$  Å, space group  $P2_12_12_1$ ,  $D_c = 1.12$ ,  $D_m = 1.12$  g cm $^{-3}$  (in  $ZnCl_2$ ),  $Z = 4$ . The diffraction intensities were collected in the  $\omega$  scan mode using graphite-monochromated Mo- $K_\alpha$

radiation on a Syntex R3 automatic diffractometer. 2235 independent reflections were measured, and 1200 were judged to be observed after correction for Lorentz, polarization, and background effects. The structure was solved by direct methods using MULTAN in the Syntex XTL program. Refinement by full-matrix least-squares, using anisotropic thermal parameters for the non-hydrogen atoms and isotropic ones for the hydrogen atoms excluding those of the five methyl groups, converged to a current  $R$ -value of 0.067. The structure, excluding the absolute configuration, of the benzoate (6) is shown in the Figure. By analysis of the c.d. spectrum ( $[\theta]_{285} +11,750$  and  $[\theta]_{213} +18,700$  in MeOH;  $[\theta]_{292} +9,620$  and  $[\theta]_{202} +21,900$  in iso-octane) of the cyclopropyl ketone (4),<sup>5</sup> the absolute configuration was then established as (2).

The alcohol (2) contains a novel fused 3,6,6,5-tetracyclic ring system in the *cis-trans-anti-trans*-configuration, and the hydroxy group is *cis* to the cyclopropane ring.

When the cyclopropyl alcohol (2) was treated with acid, it was converted in good yield into the homoallyl alcohol (7),  $C_{20}H_{34}O$ , m.p. 153–154 °C,  $[\alpha]_D +17^\circ$ , which was identical with (+)-homoverrucosan-5 $\beta$ -ol obtained by the same acid treatment of (–)-verrucosanol (1).<sup>3</sup>

We thank Dr. Y. Kushi, Hiroshima University, for the X-ray analysis and Dr. K. Kuriyama, Shionogi Research Laboratory for determination of the c.d. spectrum.

(Received, 1st May 1980; Com. 456.)

§ The atomic co-ordinates for this work are available on request from the Director of the Cambridge Crystallographic Data Centre, University Chemical Laboratory, Lensfield Rd., Cambridge CB2 1EW. Any request should be accompanied by the full literature citation for this communication.

<sup>1</sup> A. Matsuo, H. Nozaki, M. Nakayama, S. Hayashi, and D. Takaoka, *J. Chem. Soc., Chem. Commun.*, 1978, 198.

<sup>2</sup> S. Hayashi, A. Matsuo, H. Nozaki, M. Nakayama, D. Takaoka, and M. Hiroi, *Chem. Lett.*, 1978, 953.

<sup>3</sup> D. Takaoka, *J. Chem. Soc., Perkin Trans. 1*, 1979, 2711.

<sup>4</sup> H. Nozaki, A. Matsuo, Y. Kushi, M. Nakayama, S. Hayashi, D. Takaoka, and N. Kamijo, *J. Chem. Soc., Perkin Trans. 2*, 1980, 763.

<sup>5</sup> K. Kuriyama, H. Tada, Y. K. Sawa, S. Itô, and I. Itoh, *Tetrahedron Lett.*, 1968, 2539.